


# CHEMISTRY



## Important Questions

### Multiple Choice questions-

Question 1. Which of the following units is useful in relating concentration of solution with its vapour pressure?

- (a) mole fraction
- (b) parts per million
- (c) mass percentage
- (d) molality

Question 2. On dissolving sugar in water at room temperature, solution feels cool to touch. Under which of the following cases dissolution of sugar will be most rapid?

- (a) Sugar crystals in cold water.
- (b) Sugar crystals in hot water.
- (c) Powdered sugar in cold water.
- (d) Powdered sugar in hot water.

Question 3. At equilibrium the rate of dissolution of a solid solute in a volatile liquid solvent is

- (a) less than the rate of crystallisation
- (b) greater than the rate of crystallisation
- (c) equal to the rate of crystallisation
- (d) zero

Question 4. A beaker contains a solution of substance 'A'. Precipitation of substance 'A' takes place when small amount of 'A' is added to the solution. The solution is .....

- (a) saturated
- (b) supersaturated

- (c) unsaturated
- (d) concentrated

Question 5. Maximum amount of a solid solute that can be dissolved in a specified amount of a given liquid solvent does not depend upon .

- (a) temperature
- (b) nature of solute
- (c) pressure
- (d) nature of solvent

Question 6. Low concentration of oxygen in the blood and tissues of people living at high altitude is due to

- (a) low temperature
- (b) low atmospheric pressure
- (c) high atmospheric pressure
- (d) both low temperature and high atmospheric pressure

Question 7. Considering the formation, breaking and strength of hydrogen bond, predict which of the following mixtures will show a positive deviation from Raoult's law?

- (a) Methanol and acetone.
- (b) Chloroform and acetone.
- (c) Nitric acid and water.
- (d) Phenol and aniline.

Question 8. Colligative properties depend on

- (a) the nature of the solute particles dissolved in solution.
- (b) the number of solute particles in solution.

- (c) the physical properties of the solute particles dissolved in solution.
- (d) the nature of solvent particles.

Question 9. Which of the following aqueous solutions should have the highest boiling point?

- (o) 1.0 M NaOH
- (b) 1.0 M Na<sub>2</sub>SO<sub>4</sub>
- (c) 1.0 M NH<sub>4</sub>NO<sub>3</sub>
- (d) 1.0 M KNO<sub>3</sub>

Question 10. The unit of ebullioscopic constant is

- (a) K kg mol<sup>-1</sup> or K (molality)<sup>-1</sup>
- (b) mol kg K<sup>-1</sup> or K<sup>-1</sup> (molality)
- (c) kg mol<sup>-1</sup> K<sup>-1</sup> or K<sup>-1</sup> (molality)<sup>-1</sup>
- (d) K mol kg<sup>-1</sup> or K (molality)

Question 11. In comparison to a 0.01 M solution of glucose, the depression in freezing point of a 0.01 M MgCl<sub>2</sub> solution is .....

- (a) the same
- (b) about twice
- (c) about three times
- (d) about six times

Question 12. An unripe mango placed in a concentrated salt solution to prepare pickle shrivels because .....

- (a) it gains water due to osmosis.
- (b) it loses water due to reverse osmosis.
- (c) it gains water due to reverse osmosis.

(d) it loses water due to osmosis.

Question 13. At a given temperature, osmotic pressure of a concentrated solution of a substance

(a) is higher than that of a dilute solution.

(b) is lower than that of a dilute solution.

(c) is same as that of a dilute solution.

(d) cannot be compared with osmotic pressure of dilute solution.

Question 14. Which of the following statements is false?

(a) Two different solutions of sucrose of same molality prepared in different solvents will have the same depression in freezing point.

(b) The osmotic pressure of a solution is given by the equation  $\pi = cRT$  (where  $c$  is the molarity of the solution).

(c) Decreasing order of osmotic pressure for 0.01 M aqueous solutions of barium chloride, potassium chloride, acetic acid and sucrose is  $BaCl_2 > KCl > CH_3COOH > \text{sucrose}$ .

(d) According to Raoult's law, the vapour pressure exerted by a volatile component of a solution is directly proportional to its mole fraction in the solution.

Question 15. The values of Van't Hoff factors for KCl, NaCl and  $K_2SO_4$ , respectively, are

(a) 2, 2 and 2

(b) 2, 2 and 3

(c) 1, 1 and 2

(d) 1, 1 and 1

### Very Short Question:

Question 1. Define the term – solubility?

Question 2. What is the effect of pressure on solubility of a gas?

Question 3. State Henry's Law.

Question 4. State Raoult's Law.

Question 5. What are the factors on which vapour pressure depends?

Question 6. The vapour pressure of solvent gets lowered, when a non-volatile solute is added to it. Why?

Question 7. Name two ways by which vapour pressure of a liquid can be lowered.

Question 8. Define solution?

Question 9. Define the following terms:

(a) Molality

(b) Molarity

Question 10. How does change in temperature changes the molarity and molality values?

### Short Questions:

Question 1 Find the molality and molarity of a 15% solution of  $\text{H}_2\text{SO}_4$  when its density is

a)  $1.10 \text{ g/cm}^3$  & molar mass = 98 amu.

Question 2. Calculate the mole fraction of ethanol and water in a sample of rectified spirit which contains 46% ethanol by mass?

Question 3. Calculate the % composition in terms of mass of a solution obtained by mixing 300g of a 25% & 400 g of a 40% solution by mass?

Question 4. One litre of sea water weight 1030g and contains about  $6 \times 10^{-3} \text{ g}$  of dissolved  $\text{O}_2$ . Calculate the concentration of dissolved oxygen in ppm?

Question 5. The density of 85% phosphoric acid is.  $1.70 \text{ g/cm}^3$  What is the volume of a solution that contains 17g of phosphoric acid?

Question 6. Define the term azeotrope?

Question 7. Obtain a relationship between relative lowering of vapour pressure and mole fraction of solute?

Question 8. Draw the graphs of both deviations from ideal behaviours?

Question 9. A weak electrolyte AB is 5% dissociated in aqueous solution? What is the

freezing point of a 0.10 molar aqueous solution of AB?  $K_f = 1.86 \text{ deg/molal}$ ?

Question 10. Henry's law constant for the molality of methane in benzene at 298 K is  $4.27 \times 10^5 \text{ mmHg}$ . Calculate the solubility of methane in benzene at 298 K under 760 mm Hg.

### Long Questions:

#### Question 1.

The vapour pressure of  $\text{CS}_2$  at  $500^\circ\text{C}$  is 854 mm Hg. A solution of 2.0g sulphur in 100g of  $\text{CS}_2$  has a vapour pressure of 848.9 mm Hg. Calculate the formula of sulphur molecule.

**Question 2.** Calculate the mass percentage of benzene ( $\text{C}_6\text{H}_6$ ) and carbon tetrachloride ( $\text{CCl}_4$ ) if 22 g of benzene is dissolved in 122 g of carbon tetrachloride.

**Question 3.** Calculate the mole fraction of benzene in solution containing 30% by mass in carbon tetrachloride.

**Question 4.** Calculate the molarity of each of the following solutions: (a) 30 g of  $\text{CO}(\text{NO}_3)_2 \cdot 6\text{H}_2\text{O}$  in 4.3 L of solution (b) 30 mL of 0.5 M  $\text{H}_2\text{SO}_4$  diluted to 500mL.

**Question 5.** Calculate (a) molality (b) molarity and (c) mole fraction of KI if the density of 20% (mass/mass) aqueous KI is  $1.202 \text{ gml}^{-1}$ .

### Assertion and Reason Questions:

1. In these questions, a statement of assertion followed by a statement of reason is given. Choose the correct answer out of the following choices.

- Assertion and reason both are correct statements and reason is correct explanation for assertion.
- Assertion and reason both are correct statements but reason is not correct explanation for assertion.
- Assertion is correct statement but reason is wrong statement.
- Assertion is wrong statement but reason is correct statement.

**Assertion:** Camphor is used as a solvent in the determination of molecular masses of naphthalene, anthracene, etc.

**Reason:** Camphor has high molal elevation constant.

2. In these questions, a statement of assertion followed by a statement of reason is given. Choose the correct answer out of the following choices.

- a) Assertion and reason both are correct statements and reason is correct explanation for assertion.
- b) Assertion and reason both are correct statements but reason is not correct explanation for assertion.
- c) Assertion is correct statement but reason is wrong statement.
- d) Assertion is wrong statement but reason is correct statement.

**Assertion:** Reverse osmosis is used in the desalination of sea water.

**Reason:** When pressure more than osmotic pressure is applied, pure water is squeezed out of the sea water through the membrane.

### Case Study Questions:

1. Read the passage given below and answer the following questions:

The solubility of gases increases with increase of pressure. William Henry made a systematic investigation of the solubility of a gas in a liquid. According to Henry's law "the mass of a gas dissolved per unit volume of the solvent at constant temperature is directly proportional to the pressure of the gas in equilibrium with the solution". Dalton during the same period also concluded independently that the solubility of a gas in a liquid solution depends upon the partial pressure of the gas. If we use the mole fraction of gas in the solution as a measure of its solubility, then Henry's law can be modified as "the partial pressure of the gas in the vapour phase is directly proportional to the mole fraction of the gas in the solution"

The following questions are multiple choice questions. Choose the most appropriate answer:

- (i) Henry's law constant for the solubility of methane in benzene at 298K is  $4.27 \times 10^5$  mm Hg. The solubility of methane in benzene at 298K under 760mm Hg is:
  - a)  $4.27 \times 10^{-5}$
  - b)  $1.78 \times 10^{-3}$
  - c)  $4.27 \times 10^{-3}$
  - d)  $1.78 \times 10^{-5}$
  
- (ii) The partial pressure of ethane over a saturated solution containing  $6.56 \times 10^{-2}$ g of ethane is 1 bar. If the solution contains  $5.00 \times 10^{-2}$ g of ethane then what will be the partial pressure (in bar) of the gas?
  - a) 0.762
  - b) 1.312
  - c) 3.81
  - d) 5.0

(iii)  $K_H$  (K bar) values for  $Ar_{(g)}$ ,  $CO_{2(g)}$ ,  $HCHO_{(g)}$  and  $CH_{4(g)}$  are 40.39, 1.67,  $1.83 \times 10^{-5}$  and 0.413 respectively. Arrange these gases in the order of their increasing solubility. Arrange these gases in the order of their increasing solubility.

- a)  $HCHO < CH_4 < CO_2 < Ar$
- b)  $HCHO < CO_2 < CH_4 < Ar$
- c)  $Ar < CO_2 < CH_4 < HCHO$
- d)  $Ar < CH_4 < CO_2 < HCHO$

(iv) When a gas is bubbled through water at 298K, a very dilute solution of the gas is obtained. Henry's law constant for the gas at 298K is 150 kbar. If the gas exerts a partial pressure of 2 bar, the number of millimoles of the gas dissolved in 1L of water is:

- a) 0.55
- b) 0.87
- c) 0.37
- d) 0.66

(v) Which of the following statements is correct?

- a)  $K_H$  increases with increase of temperature.
- b)  $K_H$  decreases with increase of temperature.
- c)  $K_H$  remains constant with increase of temperature.
- d)  $K_H$  first increases then decreases, with increase of temperature.

2. Read the passage given below and answer the following questions

Few colligative properties are:

1. Relative lowering of vapour pressure: depends only on molar concentration of solute (mole fraction) and independent of its nature.
2. Depression in freezing point: it is proportional to the molal concentration of solution.
3. Elevation of boiling point: it is proportional to the molal concentration of solute.
4. Osmotic pressure: it is proportional to the molar concentration of solute

A solution of glucose is prepared with 0.052 g of glucose in 80.2 g of water. ( $K_f = 1.86K \text{ kg mol}^{-1}$  and  $K_b = 5.2K \text{ kg mol}^{-1}$ )

The following questions are multiple choice questions. Choose the most appropriate answer:

(i) Molality of the given solution is.

- a) 0.0052m

- b) 0.0036m
- c) 0.0006m
- d) 1.29m

(ii) Boiling point for the solution will be.

- a) 373.05K
- b) 373.15K
- c) 373.02K
- d) 373.02K

(iii) The depression in freezing point of solution will be.

- a) 0.0187K
- b) 0.035K
- c) 0.082K
- d) 0.067K

(iv) Mole fraction of glucose in the given solution is.

- a)  $6.28 \times 10^{-5}$
- b)  $6.28 \times 10^{-4}$
- c) 0.00625
- d) 0.00028

(v) If same amount of sucrose ( $C_{12}H_{22}O_{11}$ ) is taken instead of glucose, then.

- a) Elevation in boiling point will be higher.
- b) Depression in freezing point will be higher.
- c) Depression in freezing point will be lower.
- d) Both (a) and (b).

### Answers key

#### MCQ answers:

1. Answer: (a) mole fraction
2. Answer: (d) Powdered sugar in hot water.
3. Answer: (c) equal to the rate of crystallisation
4. Answer: (b) supersaturated

5. Answer: (c) pressure
6. Answer: (b) low atmospheric pressure
7. Answer: (a) Methanol and acetone.
8. Answer: (b) the number of solute particles in solution.
9. Answer: (b) 1.0 M Na<sub>2</sub>SO<sub>4</sub>
10. Answer: (a) K kg mol<sup>-1</sup> or K (molality)<sup>-1</sup>
11. Answer: (c) about three times
12. Answer: (d) it loses water due to osmosis.
13. Answer: (a) is higher than that of a dilute solution.
14. Answer: (a) Two different solutions of sucrose of same molality prepared in different solvents will have the same depression in freezing point.
15. Answer: (b) 2, 2 and 3

#### Very Short Answers:

1. The maximum amount of a substance that can be dissolved in a specified amount of solvent is called its solubility.
2. The solubility of a gas increases with increases of pressure.
3. Answer: Henry's Law states that at a constant temperature the solubility of a gas in a liquid is directly proportional to the pressure of the gas.
4. Answer: Raoult's Law states that for a solution of volatile liquids, the partial vapour pressure of each component in the solution is directly proportional to its mole fraction.
5. Answer: The factors on which vapour pressure depends are –  
1) Temperature of the liquid.    2) Nature of the liquid.
6. Answer: When a non-volatile solute is added to a solvent, the surface area for escape of solvent molecules decreases and vapour pressure gets lowered.
7. Answer: The two ways by which vapour pressure can be lowered are –  
1) By decreasing the temperature.

- 2) By adding a non- volatile solute.
8. Solutions are homogeneous mixtures of two or more than two components..
9. (a) Molality is defined as the number of moles of the solute per kilogram of solvent.

$$\text{Molality (m)} = \frac{\text{Moles of solutes}}{\text{Mass of solvent in Kg}}$$

(b) Molarity (M) = Number of moles of solute dissolved in one litre of solution.

$$\text{Molarity (M)} = \frac{\text{No. of Moles of solute}}{\text{Volume of solution in litre}}$$

10. As the temperature increases, volume increases and molarity decreases whereas molality does not change with any change in temperature.

### Short Answers:

#### 1. Answer

$$\text{Volume} = \text{mass/density}$$

$$= 100 \text{ g} / 1.10 \text{ g/cm}^3 = 90.9 \text{ cm}^3$$

$$\text{Molarity} = \frac{\text{no. of moles of H}_2\text{SO}_4}{\text{volume of solution}} \times 1000$$

$$= \frac{(15/98)}{90.9} \times 1000 = 1.68 \text{ m}$$

$$\text{Molality} = \frac{\text{no. of moles of solute}}{\text{mass of solvent in kg}}$$

$$\frac{(15/98) \times 1000}{85 \text{ g}}$$

$$= 1.8 \text{ M.}$$

#### 2. Answer:

$$\text{Mass of ethanol} = 46 \text{ g}$$

$$\text{Mass of water} = 100 - 46 = 54 \text{ g}$$

Mole fraction of ethanol,

$$\frac{\frac{46}{46}}{\frac{46}{46} + \frac{54}{18}} = \frac{1}{1+3} = \frac{1}{4} = 0.25.$$

Mole fraction of water =  $1 - 0.25 = 0.75$

Mole fraction of water =  $1 - 0.25 = 0.75$

### 3. Answer:

mass of solute in 400g of 40%

$$= \frac{40}{100} \times 400 = 160\text{g}$$

Total mass of solute =  $160 + 75 = 235\text{g}$

Total mass of solution =  $400 + 300 = 700\text{g}$

$$\text{Mass\% of solute} = \frac{\text{mass of solute}}{\text{Total mass of solution}} \times 100$$

$$= \frac{235}{700} \times 100 = 33.57\%$$

Mass % of solvent =  $100 - 33.57 = 66.43\%$

### 4. Answer:

mass of  $O_2 = 6 \times 10^{-3} \text{g}$

ppm of  $O_2$  in 1030 g sea water =  $\frac{\text{mass of } O_2}{\text{mass of sea water}} \times 10^6$

$$= \frac{6 \times 10^{-3}}{1030} \times 10^6 = 5.8 \text{ ppm.}$$

### 5. Answer :

Ans. 85g phosphoric acid is present in 100g of solution.

17g of phosphoric acid is present in

$$\frac{100}{85} \times 17 = 20\text{g of solution}$$

$$\begin{aligned} \text{Volume of 17g of 85\% acid} &= \frac{\text{mass}}{\text{density}} \\ &= \frac{20\text{g}}{1.70\text{g/cm}^3} = 11.8\text{ cm}^3 \end{aligned}$$

**6. Answer**

A solution at certain concentration when continues to boil at constant temperature without change in its composition in solution & in vapour phase is called an azeotrope.

**7. Answer:**

According to Raoult's Law –

$$P_1 = x_1 P_1^0$$

$$\Delta P = P_1^0 - P = P_1^0 - P_1^0 X_1$$


$$= \Delta P_1 = P_1^0 (1 - X_1)$$


$$\Delta P_1 = P_1^0 X_2 \quad (X_1 + X_2 = 1)$$

$$\frac{\Delta P_1}{P_1^0} = X_2$$


$$= \frac{P_1^0 - P_1}{P_1^0} = X_2$$

Relative lowering of vapour pressure.

**8. Answer:**

**9. Answer :**

$$\text{Degree of dissociation } \alpha \text{ of AB} = \frac{.5}{100} = 0.05$$



$$\text{M} \quad 0 \quad 0$$

No. of moles dissolved

No. of moles after dissociations

$$m(1-\alpha) \quad m\alpha \quad m\alpha$$

$$0.1(1-0.05) \quad 0.1 \times 0.05 \quad 0.1 \times 0.05$$

$$\text{Total moles} = 0.1(1-0.05) + (0.1 \times 0.05) + (0.1 \times 0.05)$$

$$= 0.095 + 0.005 + 0.005 = 0.105m$$

$$\Delta T_f = K_f \cdot m$$

$$= 1.86 \times 0.105$$

$$= 0.1953 \text{ deg.}$$

$$T_f = 0^\circ\text{C} - 0.1953 = 0.1953^\circ\text{C}$$

**10. Answer :**

$$p = 760 \text{ mm Hg}$$

$$k_H = 4.27 \times 10^5 \text{ mm Hg}$$

According to Henry's law,

$$p = k_H x$$

$$x = \frac{p}{k_H}$$

$$= \frac{760 \text{ mm Hg}}{4.27 \times 10^5 \text{ mm Hg}}$$

$$= 177.99 \times 10^{-5}$$

$$= 178 \times 10^{-5} \text{ (approximately)}$$

Hence, the mole fraction of methane in benzene is  $178 \times 10^{-5}$ .

## Long Answers:

### 1. Answer:

$$P_A^\circ = 854 \text{ mm } P_A = 848.9 \text{ mm}, W_B = 2.0 \text{ g},$$

$$W_A = 100 \text{ g } M_B = ?$$

$$M_A = 12 + 2(32) = 76 \text{ g/mol (CS}_2\text{)}$$

$$\frac{P_A^\circ - P_A}{P_A^\circ} = X_B = \frac{W_B / M_B}{W_A / M_A}$$

$$M_B = W_B \times \frac{W_A}{M_A} \times \frac{P_A^\circ}{P_A^\circ - P_A}$$

$$= 2 \times \frac{100}{76} \times \frac{854}{854 - 848.9}$$

$$= 254.5 \text{ g/mol.}$$

Let the formula =  $S_x$

$$X \times 32 = 254.5 \text{ g/mol}$$

$$X = \frac{254.5}{32}$$

$$= 7.95$$

$$= 8.$$

$$= \text{Formula} = S_8$$

### 2. Answer:

$$\text{Mass percentage } C_6H_6 = \frac{\text{Mass of } C_6H_6}{\text{Total mass of the solution}} \times 100\% \text{ of}$$

$$= \frac{\text{Mass of } C_6H_6}{\text{Mass of } C_6H_6 + \text{Mass of } CCl_4} \times 100\%$$

$$= \frac{22}{22 + 122} \times 100\%$$

$$= 15.28\%$$



$$\begin{aligned} \text{Mass percentage of } \text{CCl}_4 &= \frac{\text{Mass of } \text{CCl}_4}{\text{Total mass of the solution}} \times 100\% \\ &= \frac{\text{Mass of } \text{CCl}_4}{\text{Mass of } \text{C}_6\text{H}_6 + \text{Mass of } \text{CCl}_4} \\ &= \frac{122}{22 + 122} \times 100\% \end{aligned}$$

$$= 84.72\%$$

Alternatively,

$$\text{Mass percentage of } \text{CCl}_4 = (100 - 15.28)\%$$

$$= 84.72\%$$

### 3. Answer:

Let the total mass of the solution be 100 g and the mass of benzene be 30g.

$$\begin{aligned} \therefore \text{Mass of carbon tetrachloride} &= (100 - 30) \text{ g} \\ &= 70 \text{ g} \end{aligned}$$

$$\begin{aligned} \text{Molar mass of benzene (} \text{C}_6\text{H}_6 \text{)} &= (6 \times 12 + 6 \times 1) \text{ g mol}^{-1} \\ &= 78 \text{ g mol}^{-1} \end{aligned}$$

$$\therefore \text{Number of moles of } \text{C}_6\text{H}_6 = \frac{30}{78} \text{ mol}$$

$$= 0.3846 \text{ mol}$$

$$\begin{aligned} \text{Molar mass of carbon tetrachloride (} \text{CCl}_4 \text{)} &= 1 \times 12 + 4 \times 35.5 \\ &= 154 \text{ g mol}^{-1} \end{aligned}$$

$$\therefore \text{Number of moles of } \text{CCl}_4 = \frac{70}{154} \text{ mol}$$

$$= 0.4545 \text{ mol}$$

Thus, the mole fraction of  $\text{C}_6\text{H}_6$  is given as:

$$\frac{\text{Number of moles of } C_6H_6}{\text{Number of moles of } C_6H_6 + \text{Number of moles of } CCl_4}$$

$$= \frac{0.3846}{0.3846 + 0.4545}$$

$$= 0.458$$

#### 4. Answer:

Molarity is given by:

$$\text{Molarity} = \frac{\text{Moles of solute}}{\text{Volume solution in litre}}$$

(a) Molar mass of  $CO(NO_3)_2 \cdot 6H_2O$   $59 + 2(14 + 3 \times 16) + 6 \times 18$   
 $= 291 \text{ g mol}^{-1}$

$$\text{Therefore, Moles of } CO(NO_3)_2 \cdot 6H_2O = \frac{30}{291} \text{ mol}$$

$$= 0.103 \text{ mol}$$

$$\text{Therefore, } = \frac{0.103 \text{ mol}}{4.3 \text{ L}} \text{ molarity}$$

$$= 0.023 \text{ M}$$

(b) Number of moles present in 1000 mL of 0.5  $H_2SO_4 = 0.5 \text{ mol M}$

$$\therefore \text{Number of moles present in 30 mL of 0.5 M } H_2SO_4 = \frac{0.5 \times 30}{1000} \text{ mol}$$

$$= 0.015 \text{ mol}$$

$$\text{Therefore, molarity} = \frac{0.015}{0.5 \text{ L}} \text{ mol} = 0.03 \text{ M}$$

#### 5. Answer:

(a) Molar mass of KI  $= 39 + 127 = 166 \text{ g mol}^{-1}$

20% (mass/mass) aqueous solution of KI means 20 g of KI is present in 100 g of solution.

That is,

20 g of KI is present in  $(100 - 20)$  g of water = 80 g of water

$$\text{Therefore, molality of the solution} = \frac{\text{Moles of KI}}{\text{Mass of water in kg}}$$

$$\frac{20}{166} \text{ m}$$

$$0.08$$

$$= 1.506 \text{ m}$$

$$= 1.51 \text{ m (approximately)}$$

(b) It is given that the density of the solution =  $1.202 \text{ g ml}^{-1}$

Therefore, Volume of 100 g solution =  $\frac{\text{Mass}}{\text{Density}}$

$$\frac{100 \text{ g}}{1.202 \text{ g ml}^{-1}}$$

$$= 83.19 \text{ mL}$$

$$= 83.19 \times 10^{-3} \text{ L}$$

Therefore, molarity of the solution =  $\frac{\frac{20}{166} \text{ mol}}{83.19 \times 10^{-3} \text{ L}}$

(c) Moles of KI =  $\frac{20}{166} = 0.12 \text{ mol}$

Moles of water =  $\frac{80}{18} = 4.44 \text{ mol}$

Therefore, mole fraction of KI =  $\frac{\text{Moles of KI}}{\text{Moles of KI} + \text{Moles of water}}$   
 $= \frac{0.12}{0.12 + 4.44}$   
 $= 0.0263$

### Assertion and Reason Answers:

1. (c) Assertion is correct statement but reason is wrong statement.

#### Explanation:

Camphor has high molal depression constant.

2. (a) Assertion and reason both are correct statements and reason is correct explanation for assertion.

#### Explanation:

If a pressure larger than the osmotic pressure is applied to the solution side, the pure solvent flows out of the solution to the solvent through semi-permeable membrane and this phenomenon is called reverse osmosis.

### Case Study Answers:

1. Answer :

i. (b)  $1.78 \times 10^{-3}$

**Explanation:**

$$K_H = 4.17 \times 10^5 \text{ mm HG}$$

$$p = 760 \text{ mm Hg}$$

According to Henry's law,  $P = K_H \times X_{\text{CH}_4}$

$$X_{\text{CH}_4} = \frac{P}{K_H} = \frac{760}{4.17 \times 10^5} = 1.78 \times 10^{-3}$$

ii. (a) 0.762

**Explanation:**

According to Henry's law,  $m = K_H \times p$

$$6.56 \times 10^{-2} = K_H \times 1$$

For another case,  $5 \times 10^{-2} = 6.56 \times 10^{-2} \times p$

$$p = \frac{5 \times 10^{-2}}{6.56 \times 10^{-2}} = 0.762 \text{ bar}$$

iii. (c)  $\text{Ar} < \text{CO}_2 < \text{CH}_4 < \text{HCHO}$

**Explanation:**

Higher the value of  $K_H$  at a given pressure, the lower is the solubility of the gas.

iv. (c) 0.37

**Explanation:**

The mole fraction of the gas in solution,

$$x = \frac{p}{K_H} = \frac{1}{150 \times 10^3}$$

If  $n$  is the number of moles of gas in a solution of 1 L of water containing 55.5 mol then,

$$x = \frac{n}{n+55.5} \text{ or, } \frac{n}{55.5} = \frac{1}{150 \times 10^3}$$

$[n + 55.5 \approx 55.5, \text{ as } n \text{ is very small}]$

$$n = \frac{55.5}{150} \times 10^{-3} = 0.37 \text{ millimoles}$$

v. (a)  $K_H$  increases with increase of temperature.

2. Answer :

i. (b) 0.0036m

**Explanation:**

$$m = \frac{0.052}{180} \times \frac{1000}{80.2} = 0.0036$$

ii. (c) 373.02K

**Explanation:**

$$\Delta T_b = k_b \times m = 5.2 \times 0.0036 = 0.0187 \text{ K}$$

$$T_b = 373 + 0.0187 = 373.0187 \text{ K} \approx 373.02 \text{ K}$$

iii. (d) 0.067 K

**Explanation:**

$$\Delta T_f = k_f \times m = 1.86 \times 0.0036 = 0.067 \text{ K}$$

iv. (a)  $6.28 \times 10^{-5}$

**Explanation:**

$$\text{Moles of water} = \frac{80.2}{18} = 4.455$$

$$\text{Mole fraction of glucose} = \frac{0.00028}{4.45 + 0.00028} = 6.28 \times 10^{-5}$$

v. (c) Depression in freezing point will be lower.

**Explanation:**

Depression in freezing point or elevation in boiling point is proportional to molarity, which is proportional to number of moles. For same amount, higher the molar mass of solute, lower will be number of moles. Hence, lower will be the colligative property.